

On the monad of proper factorisation systems in categories (*)

Marco Grandis

Dipartimento di Matematica, Università di Genova, Via Dodecaneso 35, 16146-Genova, Italy

E-mail address: grandis@dima.unige.it (Marco Grandis).

Abstract. It is known that factorisation systems in categories can be viewed as unitary pseudo algebras for the monad $(-)^2$, in **Cat**. We show in this note that an analogous fact holds for *proper* (i.e., *epi-mono*) factorisation systems and a suitable quotient of the former monad, deriving from a construct introduced by P. Freyd for stable homotopy. Structural similarities of the previous monad with the path endofunctor of topological spaces are considered.

MSC: 18A32; 18C15.

Key Words: Factorisation systems, 2-monads, Eilenberg-Moore algebras, pseudo algebras.

Introduction

For a category $\mathbf{X}$, the category of morphisms $\mathcal{P}\mathbf{X} = \mathbf{X}^2$ has a natural factorisation system. So equipped, it is the free category with factorisation system, on $\mathbf{X}$.

This system induces a *proper*, or *epi-mono*, factorisation system on a quotient $\mathcal{F}\mathbf{r}\mathbf{X} = \mathbf{X}^2/\mathcal{R}$ [G3], the free category with epi-mono factorisation system on $\mathbf{X}$ (the *epi-mono completion*), that generalises the Freyd embedding of the stable homotopy category of spaces in an abelian category [Fr]. "Weak subobjects" in $\mathbf{X}$, of interest for homotopy categories, correspond to ordinary subobjects in $\mathcal{F}\mathbf{r}\mathbf{X}$; other results in [G3] concern various properties of $\mathcal{F}\mathbf{r}\mathbf{X}$ that derive from weak (co)limits of $\mathbf{X}$.

Now, the "path" endofunctor $\mathcal{P} = (-)^2$ of **Cat** has an obvious 2-monad structure (with diagonal multiplication), linked to the universal property recalled above (a pseudo adjunction); it is known, since some hints in Coppey [Co] and a full proof in Korostenski - Tholen [KT], that its pseudo algebras correspond to the factorisation systems of $\mathbf{X}$. Similarly, as stated without proof in [G3], the pseudo algebras for the induced 2-monad on $\mathcal{F}\mathbf{r}\mathbf{X}$ correspond to *proper* factorisation systems of $\mathbf{X}$; more precisely, we prove here, in Theorem 4 (ii), that there is a canonical bijection between *proper* factorisation systems in $\mathbf{X}$ and pseudo isomorphism classes of unitary pseudo $\mathcal{F}\mathbf{r}$ -algebras on $\mathbf{X}$. Similar, simpler relations hold in the *strict* case: *strict factorisation systems are monadic on categories, as well as the proper such*. Structural similarities of $\mathcal{P}$ with the topological path functor $\mathcal{P}\mathbf{X} = \mathbf{X}^{[0,1]}$ are discussed at the end (Section 5).

We shall use the same notation of [G3]. For factorisation systems, one can see Freyd - Kelly [FK], Carboni - Janelidze - Kelly - Paré [CJKP], and their references; the strict version is much less used: see [G3] and Rosebrugh-Wood [RW]. Lax $\mathcal{P}$ -algebras are studied in [RT]. General lax and pseudo algebras can be found in Street [St].

(*) Work partially supported by MURST research projects.

The author acknowledges with pleasure a suggestion of F.W. Lawvere, at the origin of this note.

1. The factorisation monad. Let $\mathbf{X}$ be any category and $\mathbf{X}^2$ its category of morphisms. An object of the latter is an $\mathbf{X}$ -map $x: X' \rightarrow X''$, which we *may* write as $\hat{x}$ when it is viewed as an object of $\mathbf{X}^2$; a morphism $f = (f', f''): \hat{x} \rightarrow \hat{y}$ is a commutative square of $\mathbf{X}$, as in the left diagram

$$(1) \quad \begin{array}{ccc} X' & \xrightarrow{f'} & Y' \\ x \downarrow & & \downarrow y \\ X'' & \xrightarrow{f''} & Y'' \end{array} \qquad \begin{array}{ccccc} X' & = & X' & \xrightarrow{f'} & Y' \\ x \downarrow & & \downarrow \bar{f} & & \downarrow y \\ X'' & \xrightarrow{\quad} & Y'' & = & Y'' \end{array}$$

and the composition is obvious. The *strict factorisation* of f , shown in the right diagram, is $f = (f', 1).(1, f'')$; note that its middle object is the *diagonal* $\bar{f} = f''x = yf'$ of the square f .

Thus, $\mathbf{X}^2$ has a canonical factorisation system (*fs* for short), where the map $f = (f', f'')$ is in E (resp. in M) iff f' (resp. f'') is an isomorphism. This system contains a canonical *strict factorisation system*, where (f', f'') is in E_0 (resp. in M_0) iff f' (resp. f'') is an identity. (As in [G3, 2.1], this means that: (i) E_0, M_0 are subcategories containing all the identities; (ii) every map u has a *strictly unique* factorisation $u = me$ with $e \in E_0, m \in M_0$. A strict *fs* (E_0, M_0) is not a *fs*, of course; but, there is a unique *fs* (E, M) containing the former, where $u = me$ is in E iff m is iso, and dually. Two strict systems are said to be *equivalent* if they span the same *fs*.)

The full embedding that identifies the object X of $\mathbf{X}$, with $\hat{1}_X$

$$(2) \quad \eta\mathbf{X}: \mathbf{X} \rightarrow \mathbf{X}^2, \qquad (f: X \rightarrow Y) \mapsto (f, f): \hat{1}_X \rightarrow \hat{1}_Y,$$

makes $\mathbf{X}^2$ the *free category with factorisation system* on $\mathbf{X}$, in the "ordinary" sense (as well as in a *strict* sense): for every functor $F: \mathbf{X} \rightarrow \mathbf{A}$ with values in a category with *fs* (resp. *strict fs*), there is an extension $G: \mathbf{X}^2 \rightarrow \mathbf{A}$ that preserves factorisations (resp. *strict* factorisations), determined up to a unique functorial isomorphism (resp. *uniquely* determined): $G(\hat{x}) = \text{Im}_{\mathbf{A}}(Fx)$. The (obvious) proof is based on the canonical factorisation of $\eta\mathbf{X}(x) = (x, 1).(1, x): \hat{1}_{X'} \rightarrow \hat{1}_{X''}$ in $\mathbf{X}^2$

$$(3) \quad \begin{array}{ccccc} X' & = & X' & \xrightarrow{x} & X'' \\ 1 \downarrow & & \downarrow x & & \downarrow 1 \\ X' & \xrightarrow{x} & X'' & = & X'' \end{array} \qquad X' \xrightarrow{(1, x)} \hat{x} \xrightarrow{(x, 1)} X''.$$

One might now expect that "factorisation systems be monadic on categories", but this is only true in a relaxed 2-dimensional sense.

First, by the strict universal property, the forgetful 2-functor $U_0: \mathbf{Fs}_0\mathbf{Cat} \rightarrow \mathbf{Cat}$ (of categories with *strict fs*) has a left 2-adjoint $F_0(\mathbf{X}) = (\mathbf{X}^2; E_0, M_0)$, and we shall see that U_0 is indeed 2-monadic: the comparison 2-functor $K_0: \mathbf{Fs}_0\mathbf{Cat} \rightarrow \mathbf{P-Alg}$ establishes an isomorphism of $\mathbf{Fs}_0\mathbf{Cat}$ with the 2-category of algebras of the associated 2-monad, $P = U_0 F_0: \mathbf{Cat} \rightarrow \mathbf{Cat}$, $P(\mathbf{X}) = \mathbf{X}^2$.

Secondly, by the "relaxed" universal property, the forgetful 2-functor $U: \mathbf{FsCat} \rightarrow \mathbf{Cat}$ (of categories with *fs*) acquires a left *pseudo* adjoint 2-functor $F(\mathbf{X}) = (\mathbf{X}^2; E, M)$: the unit $\eta: 1 \rightarrow UF$ is 2-natural, but the counit is *pseudo* natural and "ill-controlled", each component $\varepsilon_{\mathbf{A}}: (|\mathbf{A}|^2; E, M) \rightarrow \mathbf{A}$ depending on a choice of images in $\mathbf{A}$; the triangle conditions are – rather – invertible 2-cells. This

would give an ill-determined *pseudo* monad structure on $\mathbb{P} = \mathbb{U}\mathbb{F} = \mathbb{U} \circ \mathbb{F}_0$, *isomorphic* to the previous 2-monad; we will therefore settle on the latter and "by-pass" the pseudo adjunction.

In fact, the structure of the category $\mathbf{2} = \{0 \rightarrow 1\}$ as a *diagonal* comonoid (with $e: \mathbf{2} \rightarrow \mathbf{1}$, $d: \mathbf{2} \rightarrow \mathbf{2} \times \mathbf{2}$) produces a *diagonal* monad on the endofunctor $\mathbb{P} = (-)^2$ of $\mathbf{Cat}$, precisely the one we are interested in. The unit $\eta\mathbf{X} = \mathbf{X}^e: \mathbf{X} \rightarrow \mathbf{X}^2$ is the canonical embedding considered above, $\eta\mathbf{X}(\mathbf{X}) = \hat{1}_{\mathbf{X}}$. The multiplication $\mu\mathbf{X} = \mathbf{X}^d: \mathbb{P}^2\mathbf{X} \rightarrow \mathbb{P}\mathbf{X}$ is a "diagonal functor" defined on $\mathbb{P}^2\mathbf{X} = \mathbf{X}^{2 \times 2}$:

- an object of $\mathbb{P}^2\mathbf{X}$ is a morphism $\xi_0 = (a_0, b_0): x_0 \rightarrow y_0$ of $\mathbb{P}\mathbf{X}$, and a commutative square in $\mathbf{X}$ (the front square of the diagram below); $\mu\mathbf{X}(\xi_0) = d_0 = b_0 x_0 = y_0 a_0$ is the diagonal of this square;
- a morphism of $\mathbb{P}^2\mathbf{X}$ is a commutative square Ξ of $\mathbb{P}\mathbf{X}$, and a commutative cube in $\mathbf{X}$; $\mu\mathbf{X}(\Xi)$ is a diagonal square of the cube

$$(4) \quad \begin{array}{ccccc} & & X'_1 & \xrightarrow{x_1} & X''_1 \\ & \nearrow f & \downarrow a_1 & \searrow d_1 & \nearrow f'' \\ X'_0 & \xrightarrow{x_0} & X''_0 & & \\ \downarrow a_0 & \nearrow g' & \downarrow d_0 & \searrow b_0 & \nearrow g'' \\ & Y'_0 & \xrightarrow{y_0} & Y''_0 & \end{array} \quad \begin{array}{l} \Xi = ((f', g'), (f'', g'')): \xi_0 \rightarrow \xi_1, \\ \xi_i = (a_i, b_i): x_i \rightarrow y_i, \\ \mu\mathbf{X}(\Xi) = (f', g''): d_0 \rightarrow d_1; \end{array}$$

μ coincides with the multiplication coming from the strict adjunction, $\mathbb{U}_0 \varepsilon_0 \mathbb{F}_0: \mathbb{P}^2 \rightarrow \mathbb{P}$ (and *would* also coincide with the pseudo multiplication $\mathbb{U}\varepsilon\mathbb{F}$, if one might control the choice of images in $\mathbb{F}\mathbf{X}$ by its strict fs).

$\mathbb{P}$ will also be called the *factorisation monad* on $\mathbf{Cat}$, while a $\mathbb{P}$ -algebra $(\mathbf{X}, t)$ will also be called a *factorisation algebra*; it consists of a functor $t: \mathbf{X}^2 \rightarrow \mathbf{X}$ such that $t.\eta\mathbf{X} = 1_{\mathbf{X}}$, $t.\mathbb{P}t = t.\mu\mathbf{X}$.

2. The proper factorisation monad. Consider now the quotient $\mathbb{F}\mathbf{r}\mathbf{X} = \mathbf{X}^2/\mathbf{R}$, modulo the "Freyd congruence" [Fr]: two parallel $\mathbf{X}^2$ -morphisms $f = (f', f''): x \rightarrow y$ and $g = (g', g''): x \rightarrow y$ are $\mathbf{R}$ -equivalent whenever their diagonals $\bar{f}$, $\bar{g}$ coincide (cf. 1.1); the morphism of $\mathbb{F}\mathbf{r}\mathbf{X}$ represented by f will be written as $[f]$ or $[f', f'']$. As a crucial effect of this congruence, if f' is epi (resp. f'' is mono) in $\mathbf{X}$, so is $[f]$ in $\mathbb{F}\mathbf{r}\mathbf{X}$.

As in [G3], a *canonical epi* (resp. *mono*) of $\mathbb{F}\mathbf{r}\mathbf{X}$ will be a morphism which can be represented as $[1, f'']$ (resp. $[f', 1]$). Every map $[f]$ has a precise *canonical factorisation* $[f] = [f', 1].[1, f'']$, formed of a canonical epi and a canonical mono (both their diagonals being $\bar{f}$). $\mathbb{F}\mathbf{r}\mathbf{X}$ has thus a *proper* strict fs (E_0, M_0) , which spans a (proper) fs (E, M) : the map $[f]: x \rightarrow y$ belongs to E iff there is some $u: Y' \rightarrow X'$ such that $yf'u = y$ (y sees f' as a *split epi*).

The full embedding $\eta'\mathbf{X} = \mathbb{P}.\eta\mathbf{X}: \mathbf{X} \rightarrow \mathbb{F}\mathbf{r}\mathbf{X}$ takes $f: X \rightarrow Y$ to $[f, f]: \hat{1}_X \rightarrow \hat{1}_Y$; $\mathbb{F}\mathbf{r}\mathbf{X}$ is thus the *free category with proper factorisation system* on $\mathbf{X}$ [G3, 2.3], called the *Freyd completion*, or *epi-mono completion* of $\mathbf{X}$. The 2-monad structure of $\mathbb{F}\mathbf{r}$, induced by the one of $\mathbb{P}$ (by-passing again a pseudo adjunction $\mathbb{F}' \dashv \mathbb{U}'$), will be called the *proper factorisation monad* on $\mathbf{Cat}$. The unit is η' . For the multiplication $\mu'\mathbf{X}: \mathbb{F}\mathbf{r}^2\mathbf{X} \rightarrow \mathbb{F}\mathbf{r}\mathbf{X}$, note that now

- an object of $\mathbb{F}\mathbf{r}^2\mathbf{X}$ is a morphism of $\mathbb{F}\mathbf{r}\mathbf{X}$, $\xi_0 = [a_0, b_0]: x_0 \rightarrow y_0$,
- a morphism of $\mathbb{F}\mathbf{r}^2\mathbf{X}$ is an equivalence class Ξ of commutative squares of $\mathbb{F}\mathbf{r}\mathbf{X}$

$$(1) \quad \Xi = [[f', g'], [f'', g'']]: (\xi_0: x_0 \rightarrow y_0) \rightarrow (\xi_1: x_1 \rightarrow y_1),$$

and we have

$$(2) \quad \mu'X(\xi_0) = d_0, \quad \mu'X(\Xi) = [f', g'']: d_0 \rightarrow d_1;$$

in fact, the class $[f', g'']: d_0 \rightarrow d_1$ is well defined, since its diagonal $g'd_0 = g''b_0x_0$ only depends on the class $[f'', g'']$ and the object x_0 . The projection p is thus a strict morphism of monads $(P, \eta, \mu) \rightarrow (F r, \eta', \mu')$, as shown in the left diagram below (with $p_2 = F r(p).pP = pF r.P(p)$)

$$(3) \quad \begin{array}{ccccc} X & \xrightarrow{\eta} & PX & \xleftarrow{\mu} & P^2X \\ \parallel & & \downarrow p & & \downarrow p_2 \\ X & \xrightarrow{\eta'} & FrX & \xleftarrow{\mu'} & Fr^2X \end{array} \quad \begin{array}{ccccc} X & \xrightarrow{\eta'} & FrX & \xleftarrow{\mu'} & Fr^2X \\ \parallel & & \downarrow t' & & \downarrow Fr(t') \\ X & \xrightarrow{\quad} & X & \xleftarrow{t'} & FrX \end{array}$$

Moreover, any Fr -algebra $t': FrX \rightarrow X$ determines a P -algebra $t = t'p: PX \rightarrow X$, while a P -algebra $t: PX \rightarrow X$ induces a Fr -algebra $t': FrX \rightarrow X$ (with $t = t'p$) iff t is compatible with R .

3. Pseudo algebras. Actually, we want to compare the 2-category $FsCat$ (of categories with fs , functors which preserve them, and natural transformations of such functors) with the 2-category $Ps-P-Alg$ of pseudo P -algebras, always understood to be *unitary* (or *normalised*).

According to a general definition (cf. [St], §2), a (unitary) *pseudo P -algebra* (X, t, ϑ) , or *factorisation pseudo algebra*, consists of a category X , a functor t (the *structure*) and a functorial isomorphism ϑ (*pseudo associativity*), so that

$$\begin{aligned} (1) \quad & t: X^2 \rightarrow X, & t.\eta X &= 1, \\ (2) \quad & \vartheta: t.Pt \cong t.\mu X: P^2X \rightarrow X, \\ (3) \quad & \vartheta(P\eta X) = 1_t = \vartheta(\eta PX): t \rightarrow t: PX \rightarrow X, \\ (4) \quad & \vartheta(P\mu X).t(P\vartheta) = \vartheta(\mu P X).\vartheta(P^2t): t.Pt.P^2t \rightarrow t.\mu X.\mu PX: P^3X \rightarrow X, \end{aligned}$$

$$\begin{array}{ccc} P^3X & \xrightarrow{\mu^P} & P^2X \\ P^2t \downarrow & \searrow P\mu & \searrow \mu \\ P^2X & & P^2X \xrightarrow{\mu} PX \\ & \searrow P\vartheta \downarrow Pt & \searrow \vartheta \downarrow t \\ & PX & \xrightarrow{t} X \end{array} \quad \begin{array}{ccc} P^3X & \xrightarrow{\mu^P} & P^2X \\ P^2t \downarrow & \searrow Pt & \searrow \mu \\ P^2X & \xrightarrow{\mu} & PX \quad PX \\ & \searrow Pt & \searrow \vartheta \downarrow t \\ & PX & \xrightarrow{t} X \end{array}$$

but here (i.e., for P) the conditions (3), (4) follow from the rest (as proved below, 4 (A), (B)).

A *morphism* $(F, \varphi): (X, t, \vartheta) \rightarrow (Y, t', \vartheta')$ of pseudo P -algebras is a functor $F: X \rightarrow Y$ with a functorial isomorphism $\varphi: F.t \rightarrow t'.PF: PX \rightarrow Y$ satisfying the following coherence conditions (again, the second is redundant for P , cf. 4 (A), (B))

$$\begin{aligned} (5) \quad & \varphi.\eta X = 1_F: X \rightarrow Y, \\ (6) \quad & \varphi\mu X.F\vartheta = \vartheta'P^2F.t'P\varphi.\varphi Pt: F.t.P t \rightarrow t'.\mu Y.P^2F: P^2X \rightarrow Y. \end{aligned}$$

Finally, a 2-cell $\alpha: (F, \varphi) \rightarrow (G, \psi)$ is just a natural transformation $\alpha: F \rightarrow G$; it is automatically coherent (cf. 4 (B))

$$(7) \quad \psi.\alpha t = t' \mathbb{P} \alpha.\varphi: F.t \rightarrow t'.\mathbb{P} G: \mathbb{P} \mathbf{X} \rightarrow \mathbf{Y}.$$

Similarly, we have the 2-category $\mathbf{Ps-Fr-Alg}$ of *pseudo Fr-algebras*, or *proper-factorisation pseudo algebras*; these amount to pseudo $\mathbb{P}$ -algebras $(\mathbf{X}, t, \vartheta)$ where both t and ϑ are consistent with R (the consistency of ϑ being redundant, cf. 4 (D)). Again, (3), (4), (6), (7) are redundant.

4. Theorem (The comparison of factorisation algebras). (i) (Coppey-Korostenski-Tholen) With respect to the diagonal 2-monad for the endofunctor $\mathbb{P} = (-)^2$ of $\mathbf{Cat}$, there is a canonical equivalence of categories – described below – between $\mathbf{FsCat}$ and $\mathbf{Ps-P-Alg}$, which induces a bijection between fs on a category $\mathbf{X}$ and pseudo isomorphism classes of pseudo $\mathbb{P}$ -algebras on $\mathbf{X}$. In the strict situation, the canonical comparison functor $K_0: \mathbf{FsCat} \rightarrow \mathbf{P-Alg}$, between strict fs and $\mathbb{P}$ -algebras, is an isomorphism.

(ii) With respect to the 2-monad of the endofunctor $\mathbf{Fr}$, the previous equivalence induces an equivalence between categories with proper factorisation systems and pseudo $\mathbf{Fr}$ -algebras, as well as a bijection between *proper* fs on a category $\mathbf{X}$ and pseudo isomorphism classes of pseudo $\mathbf{Fr}$ -algebras on $\mathbf{X}$. The comparison functor $K'_0: \mathbf{PFsCat} \rightarrow \mathbf{Fr-Alg}$, of proper strict fs, is an isomorphism.

Proof. Part (i) is mostly proved in [KT], and we only need to complete a few points.

(A) First, there is a canonical 2-functor $L: \mathbf{Ps-P-Alg} \rightarrow \mathbf{FsCat}$. Given a (unitary) pseudo $\mathbb{P}$ -algebra $(\mathbf{X}, t, \vartheta)$, every map $x: X' \rightarrow X''$ in $\mathbf{X}$ inherits a precise *t-factorisation* through the object $t(\hat{x})$, by letting the functor t act on the canonical factorisation of $\eta_{\mathbf{X}}(x) = (x, 1).(1, x)$ in $\mathbf{X}^2$ (1.3)

$$(1) \quad \begin{array}{ccccc} X' & \xlongequal{\quad} & X' & \xrightarrow{x} & X'' \\ 1 \downarrow & & \downarrow x & & \downarrow 1 \\ X' & \xrightarrow{x} & X'' & \xlongequal{\quad} & X'' \end{array} \qquad \begin{array}{ccccc} X' & \xrightarrow{\tau(x)} & t(\hat{x}) & \xrightarrow{\tau^+(x)} & X'' \end{array}$$

$$(2) \quad \begin{aligned} \tau^-(\hat{x}) &= t(1, x): X' \rightarrow t(\hat{x}), & \tau^+(\hat{x}) &= t(x, 1): t(\hat{x}) \rightarrow X'', \\ \tau^+(\hat{x}).\tau^-(\hat{x}) &= t((x, x): X' \rightarrow X'') = t.\eta(x) = x. \end{aligned}$$

E is defined as the class of $\mathbf{X}$ -maps x such that $\tau^+(\hat{x})$ is iso; dually for M . This is indeed a fs, as proved in [KT], thm. 4.4, *without* assuming the coherence condition 3.3 (cf. the Note at the end of the paper) *nor* 3.4; the fact that these properties will be obtained in (B), from the backward procedure, shows that they are redundant. (In the strict case, a strict $\mathbb{P}$ -algebra t gives a strict fs, where E_0 contains the maps x such that $\tau^+(\hat{x})$ is an identity, and dually for M_0 .)

Given a morphism $(F, \varphi): (\mathbf{X}, t, \vartheta) \rightarrow (\mathbf{Y}, t', \vartheta')$ of pseudo $\mathbb{P}$ -algebras, the fact that the functor $F: \mathbf{X} \rightarrow \mathbf{Y}$ preserves the associated fs follows from the following diagram, commutative by the naturality of $\varphi: F.t \rightarrow t'.\mathbb{P} F$ on $(1, x): X' \rightarrow \hat{x}$, $(x, 1): \hat{x} \rightarrow X''$, $(1, y)$ and $(y, 1)$

$$(3) \quad \begin{array}{ccccc} & & \tau^- Fx & & \\ & & \xrightarrow{\quad} & & \\ & FX' & & t'(Fx)^\wedge & \xrightarrow{\tau^+ Fx} & FX'' \\ & \parallel & \searrow F\tau^-x & \nearrow \varphi_x & \searrow t'(Ff) & \parallel \\ FX' & \xrightarrow{F\tau^-x} & Ft(\hat{x}) & \xrightarrow{\quad} & FX'' & \xrightarrow{Ff''} \\ & \parallel & \downarrow & & \downarrow & \parallel \\ & FY' & \xrightarrow{F\tau^-y} & Ft(\hat{y}) & \xrightarrow{F\tau^+y} & FY'' \\ & \parallel & \nearrow F\tau^-y & \searrow \varphi_y & \nearrow F\tau^+y & \parallel \\ FY' & \xrightarrow{F\tau^-y} & Ft(\hat{y}) & \xrightarrow{F\tau^+y} & FY'' & \end{array}$$

Again, we do *not* need the condition 3.6: any natural iso φ such that $\varphi \cdot \eta \mathbf{X} = 1_F$ has this effect.

(B) Conversely, one can construct a 2-functor $\mathcal{K} : \mathbf{FsCat} \rightarrow \mathbf{Ps-P-Alg}$ depending on choice. Let $(\mathbf{X}, E, M)$ be a category with fs; for every map $x: X' \rightarrow X''$, let us *choose* one structural factorisation $x = \tau^+(x) \cdot \tau^-(x): X' \rightarrow t(x) \rightarrow X''$, respecting all identities: $1 = 1.1$ (We are *not* saying that this choice comes from a strict fs contained in (E, M)). By orthogonality, this choice determines *one* functor $t: \mathbf{X}^2 \rightarrow \mathbf{X}$ with this action on the objects and such that $\tau^-: \partial^- \rightarrow t$, $\tau^+: t \rightarrow \partial^+$ are natural transformations ($\partial^-, \partial^+: \mathbf{X}^2 \rightarrow \mathbf{X}$ being the domain and codomain functors)

$$(4) \quad \begin{array}{ccccc} X' & \xrightarrow{\tau^-x} & t(x) & \xrightarrow{\tau^+x} & X'' \\ f' \downarrow & & \downarrow t(f) & & \downarrow f'' \\ Y' & \xrightarrow{\tau^-y} & t(y) & \xrightarrow{\tau^+y} & Y'' \end{array} \quad f = (f', f''): x \rightarrow y.$$

Now $t \cdot \eta(X) = t(1_X) = X$. Moreover, let $t \cdot \mathcal{P}t$ and $t \cdot \mu \mathbf{X}: \mathcal{P}^2 \mathbf{X} \rightarrow \mathbf{X}$ operate on the object $(f', f''): x \rightarrow y$ of $\mathcal{P}^2 \mathbf{X}$, producing $t \cdot \mathcal{P}t(f', f'') = Z'$ and $t \cdot \mu \mathbf{X}(f', f'') = t(\bar{f}) = Z''$

$$(5) \quad \begin{array}{ccccc} X' & \xrightarrow{x'} & t(x) & \xrightarrow{x''} & X'' \\ f' \downarrow & & \downarrow z' & & \downarrow f'' \\ & & Z' & & \\ & & \downarrow z'' & & \\ Y' & \xrightarrow{y'} & t(y) & \xrightarrow{y''} & Y'' \end{array} \quad \begin{array}{ccccc} X' & \xrightarrow{x} & & \xrightarrow{\quad} & X'' \\ f' \downarrow & \searrow d' & & \searrow d'' & \downarrow f'' \\ & & Z'' & & \\ Y' & \xrightarrow{y} & & \xrightarrow{\quad} & Y'' \end{array}$$

so that there is precisely one isomorphism $\vartheta(f): Z' \rightarrow Z''$ linking the two EM-factorisations we have obtained for the diagonal, $\bar{f} = (y''z'').(z'x') = d'' \cdot d'$ (a strict fs would give an identity, for $\vartheta(f)$)

$$(6) \quad \vartheta(f): t \cdot \mathcal{P}t(f) \rightarrow t \cdot \mu \mathbf{X}(f), \quad y''z'' = d'' \cdot \vartheta(f), \quad \vartheta(f) \cdot (z'x') = d'.$$

The coherence relations for ϑ do hold: the first (3.3) is obvious; the second (3.4) is concerned with two natural transformations, $\vartheta(\mathcal{P}\mu \mathbf{X}) \cdot t(\mathcal{P}\vartheta)$ and $\vartheta(\mu \mathcal{P} \mathbf{X}) \cdot \vartheta(\mathcal{P}^2 t)$, that take a commutative cube $\Xi \in \text{Ob}(\mathcal{P}^3 \mathbf{X})$ to the unique isomorphism linking two precise EM-factorisations of the diagonal arrow of Ξ , through $t \cdot \mathcal{P}t \cdot \mathcal{P}^2 t(\Xi)$ and $t \cdot \mu \mathbf{X} \cdot \mu \mathcal{P} \mathbf{X}(\Xi)$, respectively.

By similar arguments, a functor $F: (\mathbf{X}, E, M) \rightarrow (\mathbf{Y}, E', M')$ that preserves fs is easily seen to produce a morphism $(F, \varphi): (\mathbf{X}, t, \vartheta) \rightarrow (\mathbf{Y}, t', \vartheta')$ of the associated pseudo $\mathcal{P}$ -algebras. Note that $\varphi: F \cdot t \rightarrow t' \cdot \mathcal{P}F: \mathcal{P} \mathbf{X} \rightarrow \mathbf{Y}$ is determined by the choices which give t and t' , and does satisfy the coherence condition 3.6, $\varphi \mu \mathbf{X} \cdot F \vartheta = \vartheta' \mathcal{P}^2 F \cdot t' \mathcal{P} \varphi \cdot \mathcal{P} \mathcal{P} t$; these two natural transformations take a commutative square $\xi \in \text{Ob}(\mathcal{P}^2 \mathbf{X})$ to the unique isomorphism linking two precise EM-factorisations of the diagonal arrow of the square, through $F \cdot t \cdot \mathcal{P}t(\xi)$ and $t' \cdot \mu \mathbf{Y} \cdot \mathcal{P}^2 F(\xi)$. Similarly, a natural transformation $\alpha: F \rightarrow G$ satisfies automatically the condition 3.7.

(C) The composite $\mathbf{FsCat} \rightarrow \mathbf{Ps-P-Alg} \rightarrow \mathbf{FsCat}$ is the identity. Let (E, M) be a fs on a category $\mathbf{X}$, (t, ϑ) the associated pseudo $\mathbf{P}$ -algebra and (E', M') the fs corresponding to the latter. Then $E' = \{x \mid \tau^+(x) \text{ is iso}\}$ plainly coincides with E , and $M' = M$.

The other composite, $\mathbf{Ps-P-Alg} \rightarrow \mathbf{FsCat} \rightarrow \mathbf{Ps-P-Alg}$, is just isomorphic to the identity. It is now sufficient to consider two pseudo $\mathbf{P}$ -algebras $(t, \vartheta), (t', \vartheta')$ on $\mathbf{X}$, giving the same factorisation system (E, M) , and prove that they are pseudo isomorphic, in a unique coherent way. Actually, for each $x: X' \rightarrow X''$ in $\mathbf{X}$ there is one iso $\varphi(x)$ linking the t - and t' -factorisation (both in (E, M))

$$(7) \quad \begin{array}{ccccc} X' & \xrightarrow{\tau^-x} & t(\hat{x}) & \xrightarrow{\tau^+x} & X'' \\ \parallel & & \downarrow \varphi^x & & \parallel \\ X' & \xrightarrow{\tau'^-x} & t'(\hat{x}) & \xrightarrow{\tau'^+x} & X'' \end{array}$$

this gives a functorial isomorphism $\varphi: t \rightarrow t': \mathbf{PX} \rightarrow \mathbf{X}$ such that $(1_X, \varphi): (X, t, \vartheta) \rightarrow (X, t', \vartheta')$ is a pseudo isomorphism of algebras.

(D) For Part (ii), we only need now to prove that, in the previous transformations, pseudo $\mathbf{F r}$ -algebras (i.e., pseudo $\mathbf{P}$ -algebras consistent with the Freyd congruence $\mathbf{R}$) correspond to proper fs.

First, the consistency of $t: \mathbf{X}^2 \rightarrow \mathbf{X}$ with $\mathbf{R}$ is sufficient to give an epi-mono factorisation system. Take, for instance, $m \in M$ (so that $u = \tau^-(m)$ is iso) and $mf_1 = h = mf_2$ in the left-hand diagram below; then, the naturality of the transformation $\tau^-: \partial^- \rightarrow t$ on the $\mathbf{R}$ -equivalent maps $(f_i, h): X' \rightarrow m$ of $\mathbf{X}^2$ gives $uf_1 = t(f_1, h) = t(f_2, h) = uf_2$ and $f_1 = f_2$

$$(8) \quad \begin{array}{ccc} X' & \xlongequal{\quad} & X' \\ f_i \downarrow & & \downarrow h \\ X & \xrightarrow{m} & Y \end{array} \qquad \begin{array}{ccc} X' & \xlongequal{\quad} & X' \\ f_i \downarrow & & \downarrow t(f_i, h) \\ X & \xrightarrow{u} & t(m) \end{array}$$

Finally, if (E, M) is epi-mono, then $t(f)$ in (3) only depends on the diagonal $\bar{f}$ of $f = (f', f'')$: $x \rightarrow y$ in $\mathbf{X}^2$, and similarly for $\vartheta(f)$ in (5). Therefore they induce a functor $t': \mathbf{F rX} \rightarrow \mathbf{X}$ and a functorial iso $\vartheta': t'.\mathbf{F r}(t') \rightarrow t'.\mu'\mathbf{X}$, which form a pseudo $\mathbf{F r}$ -algebra.

5. Remarks. A crucial tool for the proof of point (A), above, is the structure of $\mathbf{PX} = \mathbf{X}^2$ as a "path functor" (representing natural transformations): it forms a *cubical comonad* [G1, G2], well linked to the previous monad structure. This interplay already arises in the exponent category $\mathbf{2}$ – a *comonoid* and a *lattice* (more precisely, a *cubical monoid* [G1]) – and was exploited in this form in [KT], Section 1.

The cubical comonad structure, relevant for formal homotopy theory [G2], has one *degeneracy* $\eta: 1 \rightarrow \mathbf{P}$ (the previous unit), two *faces* or *co-units* $\partial^\pm: \mathbf{P} \rightarrow 1$ (domain and codomain) and two *connections* or *co-operations* $g^\pm: \mathbf{P} \rightarrow \mathbf{P}^2$

$$(1) \quad \begin{array}{ccc} X' & \xrightarrow{x} & X'' \\ x \downarrow & g^-(\hat{x}) & \parallel \\ X'' & \xlongequal{\quad} & X'' \end{array} \qquad \begin{array}{ccc} X' & \xlongequal{\quad} & X' \\ \parallel & g^+(\hat{x}) & \downarrow x \\ X' & \xrightarrow{x} & X'' \end{array}$$

(The connections have appeared above in the canonical factorisation $\eta X(x) = g^-X(\hat{x}).g^+X(\hat{x})$; the natural transformations τ^- , τ^+ can thus be obtained as $\tau^- = Pt.g^+X$, $\tau^+ = Pt.g^-X$.)

A cubical comonad satisfies axioms [G1, G2] essentially saying that ∂^ε ($\varepsilon = \pm$) is a co-unit for the corresponding connection g^ε and co-absorbant for the other, while η makes everything degenerate; moreover, the connections are co-associative. Here the two structures, monad and cubical comonad, are linked by some equations (after the coincidence of the monad-unit with the degeneracy; the last formula is actually a consequence of the co-associativity of connections):

$$(2) \quad \begin{aligned} \partial^\varepsilon \mu &= \partial^\varepsilon . P \partial^\varepsilon = \partial^\varepsilon . \partial^\varepsilon P, & \mu g^\varepsilon &= 1_{PX}, \\ P\mu . g^\varepsilon P . g^{\varepsilon'} &= \eta P, & P\mu . g^\varepsilon P . g^\varepsilon &= P\mu . P g^\varepsilon . g^\varepsilon = g^\varepsilon \quad (\varepsilon \neq \varepsilon'). \end{aligned}$$

A natural question arises – if the previous arguments have a non-trivial rebound in the usual range of homotopy, the category **Top** of topological spaces. Replace the categorical interval **2** with the topological one, $I = [0, 1]$, which is, again, a diagonal comonoid and a lattice (and an exponentiable object); thus, the path functor $PX = X^I$ is a monad and a cubical comonad, consistently as above. But here, the interest of (pseudo?) P -algebras is not clear (once we have excluded the trivial, "universal" ones: for a fixed $a \in I$, every space X has an obvious strict structure, $ev_a: PX \rightarrow X$; in the same way as each category X has two trivial P -algebras, $\partial^\pm: PX \rightarrow X$, and two trivial fs). On the other hand, one can readily note that the Kleisli category of P has for morphisms the homotopies, with "diagonal" horizontal composition: $(\beta \circ \alpha)(x; t) = \beta(\alpha(x; t); t)$, for $t \in I$.

References

- [CJKP] A. Carboni - G. Janelidze - G.M. Kelly - R. Paré, *On localization and stabilization for factorization system*, Appl. Categ. Structures **5** (1997), 1-58.
- [Co] L. Coppey, *Algèbres de décompositions et précategories*, Diagrammes **3** (1980), supplément.
- [Fr] P. Freyd, *Stable homotopy*, in: Proc. Conf. Categ. Alg., La Jolla 1965, 121-176, Springer 1966.
- [FK] P. Freyd - G.M. Kelly, *Categories of continuous functors*, J. Pure Appl. Algebra **2** (1972), 1-18.
- [G1] M. Grandis, *Cubical monads and their symmetries*, in: Proc. of the Eleventh Intern. Conf. on Topology, Trieste 1993, Rend. Ist. Mat. Univ. Trieste, **25** (1993), 223-262.
- [G2] M. Grandis, *Categorically algebraic foundations for homotopical algebra*, Appl. Categ. Structures **5** (1997), 363-413.
- [G3] M. Grandis, *Weak subobjects and the epi-monic completion of a category*, J. Pure Appl. Algebra **154** (2000), 193-212.
- [KT] M. Korostenski - W. Tholen, *Factorization systems as Eilenberg-Moore algebras*, J. Pure Appl. Algebra **85** (1993), 57-72.
- [RW] R. Rosebrugh - R.J. Wood, *Factorization systems and distributive laws*, Preprint available at: www.mta.ca/~rrosebru/
- [RT] J. Rosicky - W. Tholen, *Lax factorization algebras*, Preprint available at: www.math.yorku.ca/Who/Faculty/Tholen/
- [St] R. Street, *Fibrations and Yoneda's lemma in a 2-category*, in: Category Seminar, Sydney 1972/73, Lecture Notes in Mathematics, Vol. 420, Springer, Berlin 1974, pp. 104-133.